

*Lauterbornia* H. 25: 117-123, Dinkelscherben, Juni 1996

## On the Chironomid fauna from Bulgarian inland waters

### [Chironomidae in Bulgarischen Binnengewässern]

Stefan Stoichev

With 1 table

**Schlagwörter:** Chironomidae, Diptera, Insecta, Bulgarien, Donau, Fluß, See, Baggersee, Stausee, Dominanz

**From a total of 151 chironomid species in various Bulgarian water bodies one is new for Bulgaria. The species found are relatively estimated by the dominant analysis on the base of 801 samples.**

**Von 151 Chironomiden aus verschiedenen bulgarischen Gewässern ist 1 neu für Bulgarien. Auf Grund von 801 Proben wird die relative Abundanz durch eine Dominanzanalyse berechnet.**

### 1 Introduction

*Chironomidae* is a widely distributed family of insects. Chironomid larvae inhabit various kinds of freshwater habitats, as well as the littoral of seas and oceans. They make up a considerable proportion of the benthic biomass. The density of their populations sometimes reaches several hundred thousand individuals per square meter (PANKRATOVA 1970).

Species of the family Chironomidae are interesting from both practical and theoretical points of view. Their larvae can be used as bioindicators of water contamination (SLADECEK 1964). Chironomids are also of great interest from a general biological point of view. They are models for establishing the general principles of speciation (role of chromosome rearrangements in speciation, intraspecific differentiation) (ACTON 1958, MICHAILOVA 1985a, 1985b, 1989). These widely distributed insects of importance to hydrobiology, ecology, genetics and cytogenetics have still not been fully studied from the systematic aspect.

General information about the chironomids in Bulgaria was given by RUSSEV (1966, 1967, 1984, 1994), KOVACHEV (1986) and UZUNOV & al. (1981). Recent information on the Bulgarian chironomids was reported in more details by STOICHEV (1994), STOICHEV (in press), KOVACHEV & STOICHEV (in press). The present paper gives new information on the chironomid fauna from various types of water basins according to the quantitative presence, frequency of occurrence and the distribution in the various habitats. A dominant analysis has also been made.

## 2 Material and methods

The basic part of the investigated material from Bulgaria is a collection of Prof. Valkanov and Prof. Russev gathered 1946-1993. To the collection were added the samples of the author from the rivers Danube, Jantra, Iskar, Vit, Ogosta, Rositsa, Mesta, Maritsa, Struma, Tunja, Kamchija, Ropotamo, Veleka, Rezovska, from the lakes Shabla-Ezerets, Srebarna, from the sand pit lakes near the villages of Dolni Bogrov, Chelopechene, Chepintsi, from the reservoirs Koprinka and Stambolijski, from the high mountain lakes Bliznaka, Babreka, Okoto in Rila mountain. A total of 801 samples

### Correction

In accordance with Tab. 1 please add 1 to the number of each station on page 118: 1=2, 2=3.....36=37; 1 is omitted.

3. Vojnishka-23; 4. Ogosta-25; 10. Iskar-7; 16. Mesta-21; Ropotamo-21; 23. Rila mountain-12; 27. Okoto lake, Rila mountain-8; 28. Argirovo lake, Pirin mountain-9; 29. Georgijsko lake, Pirin mountain-9; 30. Shabla lake-28; 31. Srebarna lake-36; 32. Dolni Bogrov lake-28; 33. Chelopechene lake-28; 34. Chepintsi lake-28; 35. Koprinka reservoir-24; 36. Batak reservoir-21.

The analysis of dominant quantitative presence (frequency of occurrence pF, frequency of dominance DF, range of dominance Dt in %) was made by the method of DE VRIES (1937). The determination and the presentation of the species was made according to PANKRATOVA (1970, 1977, 1983). The research of TESKEY (1981) has also been used.

## 3 Species composition

A total of 151 species from 38 genera and 5 subfamilies are included in the present study (Tab. 1). Compared to the chironomid fauna of Germany (KRETZSCHMAR & BÖTTGER 1994), of the European part of the former Soviet Union (PANKRATOVA 1970, 1977, 1983), of Romania (CURE 1964, 1985, 1989) and of Turkey (REISS 1986) the Bulgarian chironomid fauna has been recently well investigated. *Stictochironomus pictulus* is new for the Bulgarian fauna (marked by +).

**Tab. 1: Qualitative composition, frequency of occurrence and a dominant analysis of chironomid species found (Abbreviations see chapter 2 above)**

Taxon	Station	pF %	DF %	Dt %
<b>CHIRONOMINAE</b>				
<i>Beckidia zabolotzkyyi</i> (GOETGHEBUER)	12-16	1,12		
<i>Chironomus</i> gr. <i>plumosus</i> KIEFFER	8-26, 29-37	78,52	62,42	79,49
<i>Ch. riparius</i> MEIGEN	2-37	96,25	72,40	75,22
<i>Ch. halophilus</i> KIEFFER	22-25	0,37		
<i>Chironomus salinarius</i> KIEFFER	22-25	0,24		
<i>Cryptochironomus defectus</i> KIEFFER	2-37	98,00	76,27	77,82
<i>Cryptochironomus</i> gr. <i>defectus</i> KIEFFER	11-14, 16-37	74,03	61,54	83,12
<i>Cr. fridmanae</i> TSCHERNOVSKU	11-14, 16	0,62		
<i>Cryptochironomus</i> sp.	2	0,12		
<i>Demeijerea</i> sp.	12	0,12		
<i>Demicryptochironomus vulneratus</i> (ZETTERSTEDT)	12-15, 17-21	0,62		
<i>Demicryptochironomus</i> sp.	2	0,12		
<i>Dicrotendipes</i> gr. <i>nervosus</i> (STAEGER)	10-18, 26	1,37		
<i>D. nervosus</i> (STAEGER)	10-17, 27, 33	1,87		
<i>D. tritonus</i> (KIEFFER)	2, 11-13	0,49		
<i>D. dubia</i> CURE	10-12	0,37		
<i>Dicrotendipes</i> sp.	22	0,12		
<i>Einfeldia pagana</i> (MEIGEN)	11, 12	0,12		
<i>Endochironomus tendens</i> FABRICIUS	10-15, 33	1,20	0,99	79,83
<i>E. dispar</i> (MEIGEN)	12-15	0,74		
<i>Endochironomus</i> sp.	2	0,12		

Taxon	Station	pF %	DF %	Dt %
<i>Glyptotendipes glaucus</i> (MEIGEN)	12, 14, 31	0,49		
<i>G. gripekovani</i> KIEFFER	11-14, 31	0,49		
<i>G. pallens</i> (MEIGEN)	2, 3, 12-15	1,49		
<i>G. caulicola</i> (KIEFFER)	31	0,12		
<i>Glyptotendipes</i> sp.	2	0,12		
<i>Microtendipes tarsalis</i> (WALKER)	12, 13	0,24		
<i>Microtendipes gr. chloris</i> (MEIGEN)	12-17	2,12	0,12	5,66
<i>M. chloris</i> (MEIGEN)	12-15	1,12		
<i>M. pedellus</i> (DE GEER)	12-14	0,24		
<i>M. viridiscutellata</i> (GOETGHEBUER)	12-14	0,37		
<i>M. junci</i> (MEIGEN)	11-13	0,12		
<i>Microtendipes</i> sp.	22	0,12		
<i>Neozavrelia fuldensis</i> FITTKAU	12	0,12		
<i>Parachironomus kuzini</i> SHILOVA	12, 13	0,24		
<i>Paratanytarsus</i> sp.	13	0,12		
<i>Paratendipes nudisquama</i> (EDWARDS)	2-17, 19, 20	4,86	0,99	20,37
<i>P. albimanus</i> MEIGEN	10-14	3,87		
<i>P. fuscimanus</i> KIEFFER	10, 11	0,24		
<i>P. transcaucasicus</i> TSCHERNOVSKIJ	17-19	0,37		
<i>P. sordens</i> (VAN DER WULP)	17, 25	0,24		
<i>Paratendipes</i> sp.	2	0,12		
<i>Polypedium scalaenum</i> (TSCHERNOVSKIJ)	2, 3, 12, 13	0,87		
<i>P. aberrans</i> TSCHERNOVSKIJ	3, 12, 13	0,37		
<i>P. gaedii</i> (MEIGEN)	10, 11, 13-15	0,99		
<i>P. convictum</i> (WALKER)	2, 12, 14	0,37		
<i>P. pedestre</i> MEIGEN	13,14	0,24		
<i>P. tetracrenatum</i> HIRVENOVA	2, 14	0,12		
<i>Polypedium gr. nubeculosum</i> (MEIGEN)	2, 11, 14	0,37		
<i>P. nubeculosum</i> (MEIGEN)	2, 12, 14, 31	0,37		
<i>P. bicrenatum</i> KIEFFER	2	0,12		
<i>Stictochironomus pictulus</i> (MEIGEN) +	29	0,12		
<i>Tanytarsus gr. gregarius</i> KIEFFER	10, 11, 13-15, 17-21, 23-27	22,84	15,10	66,11
<i>T. gregarius</i> KIEFFER	11-14, 16, 18-24	14,85		
<i>T. longipes</i> AKHRODOV	2-11	11,36		
<i>T. mancus</i> WALKER	3-7, 11-13	8,23	6,36	77,27
<i>T. sexdentatus</i> (TSCHERNOVSKIJ)	14,15	2,74	0,74	27,00
<i>T. medius</i> REISS & FITTKAU	11, 13-15, 17, 19-23	24,84	0,24	0,96
<i>T. mendax</i> KIEFFER	12	0,24		
<i>T. pallidicornis</i> (WALKER)	2	0,12		
<i>Tanytarsus</i> sp.	25	0,12		
ORTHOCLADIINAE				
<i>Acrycotopus lucens</i> (ZETTERSTEDT)	12-16, 18	21,72	15,60	71,82
<i>Brillia gr. modesta</i> (MEIGEN)	2	0,12		
<i>B. modesta</i> (MEIGEN)	2	0,37		
<i>B. longifurca</i> KIEFFER	2	0,24		
<i>Crycotopus gr. algarum</i> (KIEFFER)	2-11, 13-25, 29	62,17	39,70	63,85
<i>Cr. algarum</i> (KIEFFER)	2-8, 11, 31	37,57	6,36	16,92
<i>Cr. alpestris</i> GOETGHEBUER	2, 28	0,62	0,49	79,03
<i>Cr. curtus</i> (HIRVENOVA)	12, 16-19	1,62		
<i>Cr. fuscus</i> (KIEFFER)	12, 16, 17, 20, 21	1,12	0,87	77,67
<i>Cr. trifascia</i> EDWARDS	2	0,24		
<i>Crycotopus gr. silvestris</i> (FABRICIUS)	2, 4, 6, 7, 11-17, 19-26, 32-37	75,15	61,42	81,72
<i>Cr. silvestris</i> (FABRICIUS)	2-37	81,14	48,81	60,15
<i>Cr. bicinctus</i> (MEIGEN)	22-24	0,87		
<i>Cr. tremulus</i> (LINNÉ)	22-25	1,37		
<i>Cr. vierriensis</i> (GOETGHEBUER)	2	0,24		
<i>Cr. ornatus</i> (MEIGEN)	31, 32	0,62		
<i>Crycotopus</i> sp.	31	0,12		
<i>Eukiefferiella cf. similis</i> GOETGHEBUER	3-7, 11, 14, 15, 17, 18, 20-23	12,23	4,11	33,61
<i>E. gracei</i> (EDWARDS)	2, 3, 10, 14-17	9,61	3,87	40,27
<i>E. ilkleyensis</i> (EDWARDS)	3-6	1,74		
<i>E. tschernovskii</i> PANKRATOVA	3, 5	0,62		
<i>E. claripennis</i> LUNDBECK	3-7	0,49		
<i>E. oxoniana</i> PANKRATOVA	2	0,24		
<i>E. lobifera</i> GOETGHEBUER	22-24	0,62		
<i>E. alpestris</i> (GOETGHEBUER)	22, 24	0,74		
<i>E. clypeata</i> (KIEFFER)	2	0,37		
<i>E. longipes</i> TSCHERNOVSKIJ (sensu PANKRATOVA 1970)	3-6, 17-20	1,37	1,12	81,75
<i>E. brevicealcar</i> (KIEFFER)	2	0,24		

Taxon	Station	pF %	DF %	Dt %
<i>E. quadridentata</i> TSCHERNOVSKIJ (sensu PANKRATOVA 1970)	12, 13	0,24		
<i>Eukiefferiella</i> sp.	2	0,12		
<i>Limnophyes</i> gr. <i>prolongatus</i> KIEFFER	9, 10, 12, 16-22	2,24		
<i>L. prolongatus</i> Kieffer	15-19, 21	4,86	3,12	64,19
<i>L. septentrionalis</i> TSCHERNOVSKIJ (sensu PANKRATOVA 1970)	11-14, 16-20	2,62		
<i>L. carelicus</i> TSCHERNOVSKIJ	22-24	2,12		
<i>Limnophyes</i> sp.	2	0,27		
<i>Metricnemus</i> gr. <i>hipopetricus</i> KIEFFER	11, 12, 18-21	1,12		
<i>Orthocladius barbatus</i> KIEFFER	11, 12, 16-21, 23	2,37		
<i>O. murvanidzey</i> TSCHERNOVSKIJ	11, 12, 16-18, 25-28	1,99		
<i>Orthocladius</i> gr. <i>saxicola</i> KIEFFER	9, 10, 13-15, 19, 20, 23	3,12		
<i>O. saxicola</i> KIEFFER	9, 10, 13-15	2,24		
<i>O. thienemanni</i> KIEFFER	3-7, 32	1,74		
<i>O. frigidus</i> (ZETTERSTEDT)	11, 13-15, 19, 20	1,62		
<i>O. olivaceus</i> (KIEFFER)	11, 13-15, 20-22, 24	1,49		
<i>O. saxosus</i> (TOKUNAGA)	11-15	0,99		
<i>Orthocladius</i> sp.	2	0,12		
<i>Parartocladius nudipennis</i> KIEFFER	11, 13-15, 17, 18, 20	3,87	3,12	80,62
<i>Parakiefferiella bathophila</i> (KIEFFER)	15, 17-20, 36, 37	3,74		
<i>Psectrocladius psilopterus</i> (KIEFFER)	13, 17, 18	2,49		
<i>Reocricotopus effusus</i> (WALKER)	3-7, 18	1,37		
<i>Sinotrocladius semivirens</i> KIEFFER	11, 16-19, 25	1,99		
<i>Simposiocladius xyloptila</i> (BOTH & CURE)	15-17	1,74		
<i>Smitia septentrionalis</i> TSCHERNOVSKIJ (s. PANKRATOVA 1970)	11-15, 17-21	3,62		
<i>S. zavreli</i> FITTKAU	13, 18, 19	0,99		
<i>S. contingens</i> (WALKER)	11, 12, 16-18	1,49		
<i>S. sedula</i> KONSTANTINOV	11, 14, 16, 17	0,49		
<i>Tvetenia</i> gr. <i>calvescens</i> (EDWARDS)	2-7, 9, 11, 16-25	12,23	4,36	35,65
<i>Tv. calvescens</i> (EDWARDS)	2-9, 11	8,98	2,62	29,17
<i>Tv. bavarica</i> (GOETGHEBUER)	12-15, 18, 19	4,49	0,99	22,04
<i>Tv. discoloripes</i> (GOETGHEBUER)	3-6, 8, 9, 12, 13	2,62		
<i>Tvetenia</i> sp.	11	0,12		
<i>Thienemanniella</i> gr. <i>clavicornis</i> (KIEFFER)	2, 11-15	1,37		
<i>T. clavicornis</i> (KIEFFER)	2	0,37		
<i>T. acuticornis</i> (KIEFFER)	12-15	0,87		
<i>Thienemanniella</i> sp.	11	0,12		
TANYPODINAE				
<i>Ablabesmyia</i> gr. <i>curticalcar</i> KIEFFER	2-11, 15, 19-20, 23-24, 26-32	43,44	18,85	43,39
<i>A. curticalcar</i> KIEFFER	2-37	46,44	20,34	43,79
<i>A. longistyla</i> FITTKAU	2	0,12		
<i>A. gr.monilis</i> LINNÉ	2, 11-15	1,87		
<i>A. monilis</i> LINNÉ	2	0,24		
<i>Ablabesmyia</i> sp.	0,22	0,12		
<i>Anatopynia plumipes</i> FRIES	2, 11-15, 19-26	4,74	0,74	15,61
<i>Apsectrotanipus trifascipennis</i> (ZETTERSTEDT)	2	0,24		
<i>Clynotanipus nervosus</i> MEIGEN	13, 15, 18, 19	1,12		
<i>Clynotanipus</i> sp.	2	0,12		
<i>Macropelopia nebulosa</i> (MEIGEN)	12, 13	0,37		
<i>Macropelopia</i> sp.	12, 13	0,12		
<i>Procladius choreus</i> MEIGEN	2, 12, 13	0,99		
<i>P. ferrugineus</i> (KIEFFER)	8-15	3,62		
<i>P. nigriventris</i> KIEFFER (sensu PANKRATOVA 1978)	11	0,24		
<i>Procladius</i> sp.	31	0,12		
<i>Tanipus kratzy</i> (KIEFFER)	13,14	0,24		
<i>T. punctipennis</i> MEIGEN	10, 12-15	1,24		
<i>Thienemannimyia lentiginosa</i> FRIES	2	0,24		
<i>Zavreliemia melanura</i> MEIGEN	11,12	0,49		
<i>Z. tetrasticta</i> (KIEFFER)	11	0,12		
DIAMESINAE				
<i>Diamesa latitarsis</i> GOETGHEBUER	12	0,24		
<i>D. pseudostylata</i> TSCHERNOVSKIJ	11, 13-15	0,99		
<i>D. insignipes</i> KIEFFER	12	0,12		
PRODIAMESINAE				
<i>Prodiamesa olivacea</i> MEIGEN	12	0,24		
<i>P. rufovittata</i> GOETGHEBUER	2	0,37		

#### 4 Dominant analysis

Tab. 1 shows that the *Chironomus riparius*, *Cryptochironomus defectus*, *Crycotopus silvestris* and *Ablabesmyia curticalcar* could be found in all Bulgarian water basins. *Chironomus gr. plumosus*, *Cryptochironomus gr. defectus*, *Crycotopus gr. silvestris*, *Crycotopus gr. algarum*, *Ablabesmyia gr. curticalcar* and *Tvetenia gr. calvescens* could be found almost everywhere. The results from the dominant analysis of the species are given in Tab. 1.

According to the obtained frequency of occurrence the following classification, proposed for the first time by STOICHEV (1994), could be applied:

1. Very frequently found species (pF > 50%): *Chironomus gr. plumosus*, *Ch. riparius*, *Cryptochironomus defectus*, *Cryptochironomus gr. defectus*, *Crycotopus gr. algarum*, *C. gr. silvestris*, *C. silvestris*. Total: 7 species.
2. Frequently found species (pF 20-50%): *Tanytarsus gr. gregarius*, *T. medius*, *Acrycotopus lucens*, *Crycotopus algarum*, *Ablabesmyia gr. curticalcar*, *Ablabesmyia curticalcar*. Total: 6 species.
3. Rarely found species (pF 1-20%): *Beckidia zabolotzkyi*, *Dicrotendipes gr. nervosus*, *D. nervosus*, *Endochironomus tendens*, *Microtendipes gr. chloris*, *M. chloris*, *Paratendipes nudisquama*, *P. albimanus*, *T. gregarius*. Total: 51 species.
4. Very rare species (pF < 1%): *Chironomus chalophilus*, *Ch. salinarius*, *Cryptochironomus fridmanae*, *Dicrotendipes tritonus*, *D. dubia*, *Einfeldia pagana*, *Endochironomus dispar*. Total: 87 species.

A comparison of the index pF and the range of the dominance Dt shows that the very frequent species dominate also qualitatively in the zoobenthic complex in various water bodies in Bulgaria (Tab. 1).

Beside with the species of high pF and Dt values (*Chironomus gr. plumosus*, *Chironomus riparius*, *Cryptochironomus defectus*, *C. gr. defectus*, *C. silvestris*), species of high values of the range of dominance and of low presence and dominance frequency could be found (*Endochironomus tendens*, *Paratendipes nudisquama*, *Tanytarsus gr. gregarius*, *Acrycotopus lucens*, *Crycotopus alpestris*, *Crycotopus fuscus*, *Eukiefferiella graeci*, *E. longipes*, *Limnophyes prolongatus*, *Pararthrocladius nudipennis*; Tab. 1).

The presented data establish the stenobiontic character of some species as well. The abundant development of these species is possible only in narrow limits of the environmental conditions. Out of these limits they can not be found or they are quantitatively scanty. It is most probable that the more polluted stretches of the rivers and influenced lakes cause instability of the environment. At the places with great selfpurificational capacity a well composed and usually constant qualitative composition can be found.

## Acknowledgments

I gratefully like to thank Prof. Dr. B. Russev for the collected material; to Dr. I. Pandourski for valuable advice; to Theodora Trichkova, Nicholas Chernev and Svetoslav Cheshmedjiev for their reviewing and technical support.

Funding for this research was provided, in part, by grant from the National Fund for Scientific Research within B-433 Project.

## References

- ACTON, B. (1955): Larval group in the subgenus *Chironomus* Meigen.- Arch. Hydrobiol. **50**: 64-75, Stuttgart.
- CURE, V. (1964): Beitrage zur Kenntniss der Tendipediden (Larven) in rumanischen Donaugebiet.- Arch. Hydrobiol. **27**: 418-441, Stuttgart.
- CURE, V (1985): Chironomidae (Diptera-Nematocera) aus Rumänien unter besonderer Berücksichtigung jener aus dem hydrographischen Einzugsgebiet der Donau.- Arch. Hydrobiol. **68**: 163-217, Stuttgart.
- CURE, V. (1989): La structure spécifique et le rôle écologique de quelques populations de chironomides phytophiles (larves) associées aux macrophytes aquatiques du Danube et d'autres écosystèmes du bassin hydrographique du fleuve sur le secteur roumain.- Trav. Mus. Hist. nat. "Grigore Antipa" **30**: 267-290, Bucarest.
- DE VRIES, M. (1937): Methods used in plant sociology and agricultural botanical grassland research.- *Herbage Rev.* **5**: 76-82.
- KOVACHEV, S. & J. Uzunov (1986): Formation of macroinvertebrate communities in the course of the biological selfpurification of the Mesta River.- Arch. Hydrobiol. **72**: 427-526, Stuttgart.
- KOVACHEV, S. & S. STOICHEV (in press): The Zoobenthos from the Three Sand-Pit Lakes at different stages of their development.- *Hydrobiology* **40**, Sofia.
- KRETZSCHMAR, A. & K BÖTTGER (1994): Die Chironomidae (Diptera, Nematocera) eines kleinen, naturnahen Baches im Norddeutschen Tiefland (Kremper Au, Schleswig-Holstein).- *Lauterbornia* **19**: 161-169, Dinkelscherben.
- MICHAILOVA, P. (1985a): Tendencies in the karyotype evolution of species of the family Chironomidae (Diptera).- *Acta Zool. Bulgarica* **26**: 3-22, Sofia.
- MICHAILOVA, P. (1989): The polytene chromosomes and their significance to the systematics of the family Chironomidae (Diptera).- *Acta Zool. Fennica* **186**: 1-107, Helsinki.
- MICHAILOVA, P. (1985b): Cytological sex determination of some Chronomidae species (Diptera) (in Bulgarian, with English summary).- *National Conference in Genetics*: 100-104, Sofia.
- PANKRATOVA, V. (1970): Lichinki i kukolki komarov podsemeistva Orthocladiinae fauny SSSR (Diptera, Chironomidae=Tendipedidae).- *Opred. Faune SSSR* **102**: 1-343, Leningrad. (in Russian)
- PANKRATOVA, V (1977): Lichinki i kukolki komarov podsemeistv Podonominae i Tanypodinae fauny SSSR (Diptera, Chironomidae=Tendipedidae).- *Opred. Faune SSSR* **112**: 1-155, Leningrad. (in Russian)
- PANKRATOVA, V. (1983): Lichinki i kukolki komarov podsemeistva Chironominae fauny SSSR (Diptera, Chironomidae=Tendipedidae).- *Opred. Faune SSSR* **134**: 1-296, Leningrad. (in Russian)
- REISS, F. (1986): A contribution to the zoogeography of the Turkish Chironomidae (Diptera).- *Israel J. Entomol.* **19**: 161-170.
- RUSSEV, B. (1966): Hydrobiologische Intersuchungen der Marica.- *I. Bull. Inst. Zool. Mus. Fauna Thrakiens* **3**, Band **3**: 231-291, Sofia.

- RUSSEV, B. (1967): Hydrobiologische Untersuchungen der Marica. II. Saprobiologische Bewertung für die Jahre 1965 und 1966.- Bull. Inst. Zool. Mus. **25**: 87-99, Sofia.
- RUSSEV, B., M. NICOLOVA & M. DIMITROVA (1984): Hydrobiological and saprobiological alterations in the Tunja River.1.1955-1967.- Hydrobiology **22**: 59-73, Sofia.
- RUSSEV, B., I. YANEVA & R. DETCHEVA (1994): 3. Hydrobiologie, 3.1. Zusammensetzung der Hydrofauna, 3.1.1. Wirbellose Tiere.- In: RUSSEV, B. (Hrsg.): Limnologie der bulgarischen Donauzuflüsse: 130-255.- (Verl. "Paper Tiger") Sofia (bulg., deutsche Zsfg.).
- SLADECEK, V. (1964): Zur Ermittlung des Indikatorgewichtes in der biologischen Gewässeruntersuchung.- Arch. Hydrobiol. **60**: 241-243, Stuttgart.
- STOICHEV, S. (1994): Contribution to the study of the chironomid fauna (Diptera, Chironomidae) in the Bulgarian stretch of the Danube River and its tributaries. Frequency of occurrence and dominant analysis of the species found.- Hydrobiology **39**: 91-101, Sofia.
- STOICHEV, S. (in press): The Zoobenthos from the Lakes Shabla-Ezerets.- Hydrobiology **41**, Sofia.
- TESKEY, J. (1981): Key to the families - Larval.- In: Manual of Nearctic Diptera.I, Monogr. **27**: 1-117, (Biosystems Research Inst. Ottawa) Ottawa.
- UZUNOV, J., S. Russev, S. KOVACHEV & I. YANEVA (1981): Species Composition and Distribution of the Macrozoobenthos of the Maritsa River.- Hydrobiology **14**: 3-15, Sofia.

*Anschrift des Verfassers:* Stefan Stoichev, Bulgarian Academy of Sciences, Institute of Zoology, boul. 1 Zar Osvoboditel, 1000 Sofia, Bulgaria

*Manuskripteingang:* 27.11.1995

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Lauterbornia](#)

Jahr/Year: 1996

Band/Volume: [1996\\_25](#)

Autor(en)/Author(s): Stoichev Stefan

Artikel/Article: [On the Chironomid fauna from Bulgarian inland waters. 117-123](#)