

Six years of glacier mass balance on Mullwitzkees (Hohe Tauern) and Hallstätter Gletscher (Dachstein)

Martin Stocker-Waldhuber^{1,2}, Andrea Fischer^{1,3}, Florian Jurgeit⁴, Klaus Reingruber⁵

¹ Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Austria

² Institute for Geography, Physical Geography, Martin-Luther-University of Halle-Wittenberg, Germany

³ Institute for Meteorology and Geophysics, University of Innsbruck, Austria

⁴ Hohe Tauern National Park, Austria

⁵ BLUE SKY Wetteranalysen, Attnang, Austria

Abstract

Since 2006 the Mullwitzkees, situated within the core zone of the Hohe Tauern National Park and the Hallstätter Gletscher, situated in the Dachstein region within the northern limestone alps, are subject to mass balance monitoring programs. The mass balance of these glaciers is measured using the direct glaciological method with fixed dates by ascertaining the ice ablation at stakes and the accumulation at snow pits. The Mullwitzkees is located south of the main Alpine crest and is exposed to the south. The highest point of the glacier is the Hoher Zaun (3450 m a.s.l.) and the lowest part of the snout called Zettalunitzkees reaches down to an elevation of 2690 m. In comparison the Hallstätter Gletscher is the northernmost glacier of the Austrian Alps and is exposed to the north within an elevation range between 2150 m a.s.l. and 2900 m a.s.l., but both of the glaciers cover an area of about 3 km².

During the past six years both of the glaciers have experienced negative mass balances but at different magnitudes. Within the hydrological year 2006/2007 the specific mass balance of the Hallstätter Gletscher was about -0.3 m w.e. (water equivalent) whereas the specific balance of the Mullwitzkees was approximately five times larger (-1.4 m w.e.). Despite of this huge difference the mean specific mass balance (2006/2007 – 2011/2012) is nearly the same for both glaciers. The ELA (equilibrium line altitude) on the Mullwitzkees is about 500 m higher as it is on the Hallstätter Gletscher. These differences are mainly caused by topographic effects and different climate conditions.

Keywords

glacier, massbalance, Mullwitzkees, Venediger, Hallstätter Gletscher, Dachstein

Aims and funding of the project

The projects on Mullwitzkees and Hallstätter Gletscher aim at the measurement and interpretation of mass balance and climate data and the interpretation of the relationship of these measured parameters with respect to the current glacial recession and hydrology as well as for the development of future glacier scenarios for the both. The project on Mullwitzkees is funded by the Hohe Tauern National Park and the Hydrological Service at the Government of Tyrol. The project on Hallstätter Gletscher is funded by the Government of Upper Austria and the Energie AG. Both projects were started in 2006.

Area of study

The Mullwitzkees is situated in the Venediger Massive within the core zone of the Hohe Tauern National Park and is divided into the “innere” and “äußere” Mullwitzkees. Glacier fluctuations since the end of the LIA (Little Ice Age) are summarised by Patzelt 1973. This study focuses on the “äußere” Mullwitzkees which is therefore regarded as Mullwitzkees. The upper part of the glacier is exposed to the south and is confined by a ridge with the highest point Hoher Zaun at an elevation of 3450 m a.s.l.. The snout called Zettalunitzkees is exposed to the south-west and reaches down to an elevation of 2690 m a.s.l.. In 1998, Mullwitzkees (and Zettalunitzkees) covered an area of 3.24 km². The glacier area diminished to 3.08 km² in 2007 (STOCKER-WALDHUBER 2010).

The Hallstätter Gletscher covers an area of 3 km² and is the biggest glacier in the northern limestone Alps (LAMBRECHT & KUHN 2007). The glacier is exposed to the north and surrounded by a mountain ridge with the highest point Hoher Dachstein. From a great plateau above 2550 m a.s.l., arise three short glacier tongues. The highest point of the glacier is located at the north face of the Hoher Dachstein at an elevation of 2900 m a.s.l.. The middle tongue of the glacier reaches down to an altitude of 2150 m a.s.l.. Since the LIA maximum in 1856 the glacier area diminished by 43% (HELFRICHT 2009).




Figure 1: Distribution of the most positive (left fig.) and the most negative (right fig.) mean specific mass balance on Hallstätter Gletscher since the beginning of the measurements in 2006. The mass balance is colored gradually into 50 cm intervals within the ablation area and into 25 cm intervals within the accumulation area; the equilibrium line is plotted as a thick grey line.

The glacier boundaries in Figure 1 and Figure 2 originate from the second Austrian glacier inventory of 1998 (LAMBRECHT & KUHN 2007, KUHN et. al. 2008). For the years 2007 and 2009 the glacier's boundaries were reduced on the basis of Orthophotos and digital elevation models according to the third Austrian glacier inventory (ABERMANN et. al. 2010, STOCKER-WALDHUBER et. al. 2010). Zettalunitzkee sand Hallstätter Gletscher are also subject to measurements of glacier length by the glacier survey of the Austrian Alpine Club (e.g. PATZELT 2005, PATZELT 2006).




Figure 2: Distribution of the most positive (left fig.) and the most negative (right fig.) mean specific mass balance on Mullwitzkees since the beginning of the measurements in 2006. The mass balance is colored gradually into 50 cm intervals within the ablation area and into 25 cm intervals within the accumulation area; the equilibrium line is plotted as a thick grey line.

Method

To determine the mass balance of this glacier the direct glaciological method with fixed dates is used (HOINKES 1970). The method detects the mass gain and loss of the glacier within one year. The year is divided into the accumulation period from the 1st of October to 30th of April when a mass gain of the glacier is expected and the ablation period from the 1st of May to 30th of September when the glacier experiences a mass loss.

Ablation is measured with ablation stakes in the ablation area. During summer the free ends of the stakes are measured several times. At the 30th of April multiple snow pits are dug to measure the height and density of the accumulated snow cover and on the 30th of September this work is repeated to determine the mass gain of the glacier within the hydrological year. The direct glaciological method is described in PATERSON 1994.

Results

The Mullwitzkees (MWK) has experienced a mass loss of $4.46 \cdot 10^6 \text{ m}^3$ w.e. (water equivalent) and a specific mass balance of -1447 mm during the first year 2006/07 which is the most negative balance whereas on Hallstätter Gletscher (HSG) the mass loss devoted $0.88 \cdot 10^6 \text{ m}^3$ w.e. and a specific balance of -289 mm which was the minimum annual mass loss during the six years between 2006 and 2012. During the second year 2007/08 both glaciers got about the same negative specific balance with -642 mm (MWK) and -700 mm (HSG). In 2008/09 Hallstätter Gletscher lost nearly twice as much of its mass (-924 mm) as Mullwitzkees (-487 mm) and in 2009/10 the specific balance of Hallstätter Gletscher was -700 mm and -490 mm on Mullwitzkees. The most negative mass balance of Hallstätter Gletscher with a specific balance of -2011 mm was measured in 2010/11 which corresponds to a mass loss of about $-6 \cdot 10^6 \text{ m}^3$ w.e., whereas on Mullwitzkees the specific balance with -1303 mm was nearly the same as it was in 2006/07. In 2011/12 the specific mass balance of both of the glaciers were nearly the same with -1228 mm on Hallstätter glacier and -1276 mm on Mullwitzkees but there were huge differences between both glaciers with regard to the winter and summer balance. The specific summer mass balance of Hallstätter Gletscher in 2012 was the most negative one since the beginning of the measurements but was balanced in a large part by the most positive specific winter mass balance since 2006.

The ELA (equilibrium line altitude) was located at a mean elevation of about 2590 m a.s.l. on Hallstätter Gletscher and at 3140 m a.s.l. on Mullwitzkees during the first four years. In 2010/11 the ELA shifted up to 2822 m a.s.l. on Hallstätter Gletscher and on Mullwitzkees even above the summits and likewise during the summer 2012. Accordingly the AAR (accumulation area ratio) was very low during the year 2011/12 with 0.11 on Mullwitzkees. Due to the great amounts of winter and summer balances on Hallstätter Gletscher in 2011/12 the AAR was higher than on Mullwitzkees with 0.32. Table 1 gives an overview of the mass balance parameters of both glaciers of the years from 2006/07 to 2011/12 (FISCHER et al. 2013, STOCKER-WALDHUBER et al. 2013). Figure 1 and Figure 2 show the distribution of the most positive and most negative mean specific mass balances.

Table 1: Characteristic numbers of the mass balance on Hallstätter Gletscher and Mullwitzkees for the hydrological years from 2006/07 to 2011/12. (S: glacier area, B: total mass balance, b: mean specific mass balance, b_s: mean specific summer mass balance, b_w: mean specific winter mass balance, ELA: equilibrium line altitude, AAR: accumulation area ratio, w.e.: water equivalent)

Hallstätter Gletscher

| | mass balance | | | | | ratio | |
|-----------|-----------------|--------------------------------|---------|----------------|----------------|-------|-------|
| | S | B | b | b _s | b _w | ELA | AAR |
| | km ² | 10 ⁶ m ³ | mm w.e. | mm w.e. | mm w.e. | m | |
| 2006/2007 | 3,04 | -0,88 | -289 | -2222 | 1933 | 2581 | 0,490 |
| 2007/2008 | 3,04 | -2,13 | -700 | -3270 | 2570 | 2592 | 0,490 |
| 2008/2009 | 3,01 | -2,79 | -924 | -3069 | 2145 | 2616 | 0,341 |
| 2009/2010 | 3,016 | -2,11 | -700 | -2334 | 1634 | 2588 | 0,483 |
| 2010/2011 | 3,016 | -6,07 | -2011 | -3497 | 1486 | 2822 | 0,046 |
| 2011/2012 | 3,016 | -3,70 | -1228 | -3953 | 2725 | 2664 | 0,318 |

Mullwitzkees

| | mass balance | | | | | ratio | |
|-----------|-----------------|--------------------------------|---------|----------------|----------------|-------|-------|
| | S | B | b | b _s | b _w | ELA | AAR |
| | km ² | 10 ⁶ m ³ | mm w.e. | mm w.e. | mm w.e. | m | |
| 2006/2007 | 3,08 | -4,46 | -1447 | -2121 | 674 | 3163 | 0,207 |
| 2007/2008 | 3,08 | -1,98 | -642 | -2052 | 1410 | 3115 | 0,396 |
| 2008/2009 | 3,03 | -1,47 | -487 | -2006 | 1519 | 3116 | 0,367 |
| 2009/2010 | 3,03 | -1,48 | -490 | -1797 | 1307 | 3150 | 0,332 |
| 2010/2011 | 2,93 | -3,82 | -1303 | -2127 | 824 | **** | 0,147 |
| 2011/2012 | 2,93 | -3,74 | -1276 | -2772 | 1496 | **** | 0,111 |

Discussion

Comparing the first six years of mass balance measurements on Mullwitzkees one of the most noticeable results is the position of the accumulation area, which is displaced from the ridge to lower elevations due to wind drift during winter. The ice thickness is decreasing at the highest elevations of this glacier (SPAN et. al. 2005, FISCHER et. al. 2007). In contrast to the Mullwitzkees the mass balance of the Hallstätter Gletscher mainly depend on the snowfall amounts during winter which can be increased by orographic lifting of the Dachstein massive. Both glaciers react in a different way on the weather conditions during a specific year and benefit alternately depending on a northwesterly or southwesterly incoming flow direction.

These investigations do not relate to the status of protection, but both glaciers are located within protected areas. Nevertheless mass balance measurements are the coherency between glacier and climate and therefore it is of great importance to observe the actual conditions. These measurements help to answer questions such as how long the glaciers of the Hohe Tauern Nationalpark or within the areas of Natura 2000 tend to exist.

References

- ABERMANN, J., SEISER, B., MERAN, I., STOCKER-WALDHUBER, M., GOLLER, M. & A. FISCHER 2009/10. A new ALS glacier inventory of North Tyrol, Austria, for 2006. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 43/44, 109-119
- FISCHER, A., SPAN, N., KUHN, M. & M. BUTSCHEK 2007. Radarmessungen der Eisdicke Österreichischer Gletscher. Band II: Messungen 1999 bis 2006. *Österreichische Beiträge zu Meteorologie und Geophysik*, 39, 142pp
- FISCHER, A., STOCKER-WALDHUBER, M., REINGRUBER, K. & K. HELFRICHT 2013. Glacier mass balance of Hallstätter Gletscher, Dachstein, Austria, 2006-2012. Institute of Meteorology and Geophysics, University of Innsbruck, doi:10.1594/PANGAEA.806609
- HOINKES, H. 1970. Methoden und Möglichkeiten von Massenhaushaltsstudien auf Gletschern. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 6, 37-90
- HELFRICHT, K. 2009. Veränderungen des Massenhaushaltes am Hallstätter Gletscher seit 1856. Master's thesis, Institute for Meteorology and Geophysics, University of Innsbruck, Austria
- KUHN, M., LAMBRECHT, A., ABERMANN, J., PATZELT, G. & G. GROß 2008. Die österreichischen Gletscher 1998 und 1969, Flächen und Volumenänderungen, Austrian Academy of Sciences Press. Vienna. 123 pp.
- LAMBRECHT, A. & M. KUHN 2007. Glacier changes in the Austrian Alps during the last three decades, derived from the new Austrian glacier inventory. *Annals of Glaciology*, 46, 177-184
- PATERSON, W. 1994. *The physics of Glaciers*. Pergamon Press, 84pp
- PATZELT, G. 1973. Die neuzeitlichen Gletscherschwankungen in der Venedigergruppe (Hohe Tauern, Ostalpen) *Zeitschrift für Gletscherkunde und Glazialgeologie*, 9, 5-57
- PATZELT, G. 2005. Gletscherbericht 2003/2004. Sammelbericht über die Gletschermessungen des Oesterreichischen Alpenvereins im Jahre 2004. *Mitteilungen des Oesterreichischen Alpenvereins*. Jg. 60 (130), Heft 2/05, 24-31
- PATZELT, G. 2006. Gletscherbericht 2004/2005. Sammelbericht über die Gletschermessungen des Oesterreichischen Alpenvereins im Jahre 2005. *Bergauf* 2/2006, 6-11
- SPAN, N., FISCHER, A., KUHN, M. MASSIMO, M. & M. BUTSCHEK 2005. Radarmessungen der Eisdicke Österreichischer Gletscher. Band I: Messungen 1995 bis 1998. *Österreichische Beiträge zu Meteorologie und Geophysik*, 33, 145 pp
- STOCKER-WALDHUBER, M., WIESENEGGER, H., ABERMANN, J., HYNEK, B. & A. FISCHER 2009/10. A new glacier inventory of the province of Salzburg, Austria, 2007/2009. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 43/44, 179-186
- STOCKER-WALDHUBER, M. 2010. Untersuchung des Massenhaushalts am Mullwitzkees, im NP Hohe Tauern. Master's thesis, Institute for Meteorology and Geophysics, University of Innsbruck, Austria
- STOCKER-WALDHUBER, M., FISCHER, A. & M. KUHN 2013. Glacier mass balance Mullwitzkees, Hohe Tauern, Austria, 2006-2012. Institute of Meteorology and Geophysics, University of Innsbruck, doi:10.1594/PANGAEA.806662

Contact

Martin Stocker-Waldhuber
martin.stocker-waldhuber@oeaw.ac.at
Institute for Interdisciplinary Mountain Research
Austrian Academy of Sciences
Technikerstraße 21a
Otto-Hittmair-Platz 1
6020 Innsbruck
Austria
Tel.: +43 (0)512 507 5492

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nationalpark Hohe Tauern - Conference Volume](#)

Jahr/Year: 2013

Band/Volume: [5](#)

Autor(en)/Author(s): Stocker-Waldhuber Martin, Fischer Andrea, Jurgeit Florian, Reingruber Klaus

Artikel/Article: [Six years of glacier mass balance on Mullwitzkees \(Hohe Tauern\) and Hallstätter Gletscher \(Dachstein\) 747-750](#)