

Phyton (Austria)	Vol. 20	Fasc. 3—4	261—278	30. 9. 1980
------------------	---------	-----------	---------	-------------

The Flora of Mount Egaleo (Attica, Greece)

By

George P. SARLIS *)

With 1 figure

Received December 3, 1979

Key words: *Pteridophyta, Spermatophyta.* — Floristics, local flora. — SE Europe, Greece.

Summary

SARLIS G. P. 1980. The flora of Mount Egaleo (Attica, Greece). — Phyton (Austria) 20 (3—4): 261—278, 1 figure. — English with German summary.

Egaleo is a chain of hills in the W of Athens; the highest point reaches 468 m. The mountain as a whole is situated in the zone of Oleo-Ceratomion. At a hillside the new Botanical Garden of the University was founded. The paper contains an abstract concerning geography, geology, ecology and conditions of the vegetation. The main part is a list of 443 species of tracheophytes which have been found in the Egaleo region.

Zusammenfassung

SARLIS G. P. 1980. Die Flora des Berges Egaleo (Attika, Griechenland). — Phyton (Austria) 20 (3—4): 261—278, 1 Abbildung. — Englisch mit deutscher Zusammenfassung.

Der Egaleo ist ein Hügelzug (höchste Erhebung 468 m) im Westen von Athen, der zur Gänze im Bereich des Oleo-Ceratonions liegt. An seinem Hang wurde der neue Botanische Garten der Universität angelegt. Die Arbeit enthält einen kurzen Abriß betreffend Geographie, Geologie, Ökologie und Zustand der Vegetation. Den Hauptteil bildet eine Liste von 443 im Gebiet des Egaleo beobachteten Gefäßpflanzen.

Egaleo is situated approx. ten kilometers to the West of Athens the last wouthward extension of Mount Parnis to which it is morphologically connected. The end of the S. Egaleo range is the cap of Perama at the

*) Dr. George P. SARLIS, Department of Systematic Botany, Agricultural College of Athens, Botanikos Kipos, Iera Odos 75, Athens (301), Greece.

channel of Salamina, the southern and greatest part of which borders the continuation of this mountain (Fig. 1).

Egaleo is situated approx. ten kilometers to the West of Athens extending 17,5 km in southwestwardly direction. It is divided in two mountainous blocks by the valley driving from Daphni Cloister to Skaramangas.




Fig. 1. Map of the Egaleo region

mangas. One is the Northern Egaleo or Pikilon Mountain, whose highest point "Zaharitsa" has an altitude of 453 m above sea level and which is joined by the hill Dema to Parnis. The other is the Southern Egaleo or Mountain of Skaramangas or Korydallos, whose highest point „Pyrovolia“ reaches 468 m above sea level. The Southern and the Northern Egaleo are united by the hill of Daphni, while the western, as well as the southern side of the mountain abruptly slide down to the sea.

Northern Egaleo separates the plain of Athens from the plain of Thriasion and the valley leaves a passage, crossed by the road from Athens and Piraeus to Elefsis. This is probably the ancient passage of Pythion (the actual Daphni).

Main characteristics of Egaleo are the glens, the roughness, the aridity and the lack of vegetation, although it was woddy in ancient times (1st century A. D.), as mentioned by STATIOS (*Thibaïs XII*, 6, 620; ARRIGONI 1971).

From the geological point of view and according to the opinion of certain researchers Egaleo constitutes a "horst". The work of LEPSIUS 1893 contains a geologic map comprising a great part of the Egaleo mountain. However since the time of LEPSIUS the conceptions about the geology of Attica have changed. Studies by later researchers concerning Mount Parnis and other areas found strata of formations older than those to which the first researchers of Attica attributed the sedimentary formation.

The geologic formations that participate to the structure of Egaleo (TATARIS 1967) are most probably as follows (from the older to the younger zones):

a. A bedrock from rocks (more or less altered) of upper Palaeozoic age (Carboniferous-Permian). Within the strata of this shade-chert-sandstone formations there are intercalations of fossiliferous marly limestone with fusulines, belonging to the Uralian stage of the carboniferous period.

b. Light-gray limestones belonging to the Norian-Rhaetian stages of the Triassic-Jurassic periods.

c. Blue-gray plated marly limestones of the lower Cretaceous containing fossils of *Toucasia*, *Harpagodes* and *Requienia*.

d. Black limestones containing Rudists of the young Cretaceous subperiod.

e. Neogen marls, limestones, conglomerates and

f. Quarternaly deposits.

Serpentinized Peridotites interfere mainly between strata of the Jurassic and Cretaceous formations. As products of chemical weathering of peridotite iron-ores (mainly grained limonite) have been found at various places of Egaleo and particularly of Korydallos hill.

As far as the distribution of the aforementioned rocks in concerned, the oldest (Triassic, Jurassic, lower Cretaceous limestone, older strata) occupy the central mass of Egaleo, whereas Upper Cretaceous limestone occupy the eastern edges of it and the hill of Korydallos, as well as a narrow band by the southern borders of Egaleo, separated from the central mass by a fissure of east-western direction.

Informations concerning the flora and vegetation of Egaleo published up to date are very limited. Some of them can be found in the mimeographed study of DIAPOULIS 1968 about the Diomidios Botanical Garden in Daphni.

The mountain presents a rather poor flora and vegetation and from

that point of view it can be compared only to that of Mount Hymettus and the hill of Lycabettus (ZERLENTIS 1959, 1965).

The main characteristic of the vegetation of Egaleo is that no clear physiognomic distinction is possible between the inferior and the superior zones of the mountain. Consequently the vegetation of Egaleo belongs in its whole to the inferior zone of the sclerophyllous broad-leaved shrubs and the mediterranean conifers.

In general the vegetation is thicker and stronger on the eastern part of the mountain in comparison with the western part. The difference in thickness is due partly to the general direction of the mountain, where the western part is warmer and dryer, and partly to the intensive grazing that resulted in a deterioration and a great degradation of the natural vegetation.

In winter both parts of this mountain are exposed to strong winds, which are not favourable to high woody vegetation. Another reason is the existence of compact rocks, as well as the lack of sufficient humidity.

Finally, the lower eastern part of Egaleo has suffered a considerable influence of the anthropogenous factor up to a certain altitude.

It must be noted that on the hills of Egaleo there are untouched areas with indigenous associations of evergreen sclerophyllous broad-leaved shrubs and low-shrubby xerophiles (phrygana) belonging to the Oleo-Lentisetum of the Oleo-Ceratonion class.

The recovery of the natural vegetation of the area of the Diomidios Botanical Garden, that for many years was overgrazed by goats and sheep and was subject to the consequences of forest fires, started after the fencing -in approximately fifteen years ago.

There are old clusters of aleppo pines at the foothills mixed with cypresses. In the past different bare parts were reafforested by the Ministry of Agriculture with calabrian pine (*Pinus brutia*) and cypress (*Cupressus sp.*), as well as recently with evergreen oak (*Quercus ilex*), while at the higher points of the hills a primary vegetation of phrygana and sclerophyll bushes remained, particularly thorny and aromatic zones, that are not touched by animals. The great proportion of therophytes on the mountain Egaleo is the result of the rocky soil and the dry climate of the region.

Evidence of the vegetation existing in the past is the presence of the woody and bushy plants isolated in glens, remnants of which are still existing, as *Cupressus sempervirens*, *Pinus halepensis*, *Juniperus oxycedrus*, *J. phoenicea*, *Rubus canescens*, *Pistacia lentiscus*, *Pyrus amygdaliformis*, *Calicotome villosa*, *Anthyllis hermanniae*, *Lonicera etrusca*, *Nerium oleander*, *Quercus coccifera* etc.

Some of the plants found in Egaleo are distinguished by their preference for soils either rich or poor in calcium, constituting in a certain way an index of the soil quality. Although many of them are calcicolous pants, a few of them are growing exclusively on calcareous soils, with the result that they appear both on soils of schist and calcareous origin.

Also, a number of species is found preferentially or even exclusively on rocky grounds within the crevices of rocks or on the sides of steep rocks. Halophilous species grow near the sea, while other sensible to salt are found in the interior, reaching the peak of the mountain. The above petrophilous plants constitute interrupted associations.

We mention hereby a number of calcicolous petrophilous species of Egaleo that appear mainly on places of schist (silicous) origin:

<i>Aethionema saxatile</i>	<i>Leopoldia comosa</i>
<i>Allium ampeloprasum</i>	<i>Onobrychis ebenoides</i>
<i>Alyssum saxatile</i>	<i>Oryzopsis miliacea</i>
<i>A. montanum</i> L.	<i>Phlomis fruticosa</i>
<i>Anthyllis hermanniae</i>	<i>Psoralea bituminosa</i>
<i>Asparagus acutifolius</i>	<i>Ptilostemon chamaepeuce</i>
<i>Brassica cretica</i>	<i>Rhamnus lycioides</i> subsp. <i>graecus</i>
<i>Centranthus ruber</i>	<i>Scabiosa hymettia</i>
<i>Chrozophora tinctoria</i>	<i>Sedum hispanicum</i>
<i>Coronilla emerus</i> subsp. <i>emeroides</i>	<i>S. sediforme</i>
<i>Delphinium peregrinum</i>	<i>Stephanium tuberosus</i>
<i>Euphorbia myrsinites</i>	<i>Teucrium divaricatum</i>
<i>Galium incurvum</i>	<i>T. polium</i>
<i>Genista acanthoclada</i>	<i>Verbascum undulatum</i>
<i>Lageocia cuminoides</i>	

The following calcifuge species appear in the area of Egaleo preferentially on rocks of schist origin:

<i>Atractylis cancellata</i>	<i>Osyris alba</i>
<i>Centaurea attica</i>	<i>Phagnalon graecum</i>
<i>C. raphanina</i> subsp. <i>mixta</i>	<i>Saxifraga hederacea</i>
<i>C. solstitialis</i>	<i>Scrophularia heterophylla</i>
<i>Ceterach officinarum</i>	<i>Selaginella denticulata</i>
<i>Cheilanthes fragrans</i>	<i>Silene behen</i>
<i>Clematis cirrhosa</i>	<i>S. vulgaris</i>
<i>Cyclamen graecum</i>	<i>Umbilicus parviflorus</i>
<i>Euphorbia acanthothamnos</i>	

Finally within the restricted zone, occupied by military installations on the southern region of the mountain Egaleo, we found that the flora comprises a great number of ammophilous and halophilous species, such as:

<i>Amaranthus lividus</i>	<i>Malcolmia maritima</i>
<i>Cakile maritima</i>	<i>Medicago littoralis</i>
<i>Cynodon dactylon</i>	<i>M. marina</i>
<i>Drimia maritima</i>	<i>Oryzopsis coerulescens</i>
<i>Echium arenarium</i>	<i>Plantago coronopus</i>
<i>Glaucium flavum</i>	<i>Tribulus terrestris</i>
<i>Limonium sinuatum</i>	

The vegetation of Egaleo shows a great degradation and is in general not very different from that of the other mountains of Attica (except Mount Parnis, DIAPOULIS 1958) due to the fires, the unfair and avid exploitation of the trees, the intensive grazing, the bad treatment of excursionists, as well as the housing constructions. Already a certain number of species mentioned in the following list seem to disappear, while in older times their density must have been much greater.

If a new disturbance of the present vegetation does not follow because of the above exogenous factors, the actual low vegetation of phrygana will give its place to a new type of vegetation consisting of high bushes and low trees.

Also, because of the low altitude of the hills of Egaleo a reafforestation by mediterranean conifers (as *Pinus halepensis* and *Cupressus sempervirens*) is indicated, together with certain broad-leaved sclerophyll species, such as *Olea europaea* var. *sylvestris*, *Phillyrea latifolia*, *Quercus coccifera*, *Pistacia lentiscus* and others; this latter group present a great resistance to fire and retard its extension to the conifers especially if planted in bands.

In order to reach such an improvement of the vegetation certain protective laws must be applied by the forest authorities to avoid any further destruction of the natural environment of the region. It is most important to improve the woody vegetation of the hills and mountains within the Attic basin by all means available. Such a result cannot however be achieved without the sincere cooperation of the whole population.

List of species found in the Egaleo region

Nomenclature agrees with Flora Europaea vols. 1—4. For the species not included in this volumes, the names are in accordance with DIAPOULIS 1939—1949 or RECHINGER 1943. Instead of authors' names one of the references mentioned for nomenclature is indicated.

FAe = RECHINGER, Flora Aegaea; FE = Flora Europaea; GF = DIAPOULIS, Greece Flora. A = annual, B = biennial, P = perennial.

For many of the species mentioned the vouchers are deposited in ACA

The number of species in the Egaleo region amounts to 443, from which 178 are annuals, 23 biennials and 242 perennials (105 woody and bushy).

I. Pteridophyta

Selaginellaceae

1. *Selaginella denticulata* — P; FE 1 : 5

Sinopteridaceae

2. *Cheilanthes fragrans* — P; FE 1 : 10

Adiantaceae

3. *Adiantum capillus-veneris* — P; FE 1 : 10

Aspleniaceae

4. *Ceterach officinarum* — P; FE 1: 17

II. *Spermatophyta*

A. *Gymnospermae*

Pinaceae

5. *Pinus halepensis* — P; FE I: 35

Cupressaceae

6. *Cypressus sempervirens* — P; FE 1: 37 7. *Juniperus oxycedrus* — P; FE 1: 38
6a. *C. s. f. horizontalis* — P; FE 1: 37 8. *J. oxycedrus* subsp. *macrocarpa* — P; FE 1: 38
9. *J. phoenicea* — P; FE 1: 39

Ephedraceae

10. *Ephedra fragilis* subsp. *campylopoda* — P; FE 1: 40

B. *Angiospermae*

a. *Dicotyledones*

Fagaceae

11. *Quercus coccifera* — P; FE 1: 62 12. *Q. ilex* — P; FE 1: 62

Moraceae

13. *Ficus carica* — P; FE 1: 67

Urticaceae

14. *Urtica dioica* — P; FE 1: 68 16. *U. pilulifera* — A; FE 1: 68
15. *U. urens* — A; FE 1: 68 17. *Parietaria diffusa* — P; FE 1: 69

Santalaceae

18. *Osyris alba* — P; FE 1: 70

Aristolochiaceae

19. *Aristolochia microstoma* — P; FE 1: 74

Polygonaceae

20. *Polygonum arenarium* — A; FE 1: 78 23. *R. pulcher* — B; FE 1: 87
21. *P. aviculare* — A; FE 1: 78 24. *R. bucephalophorus* — A; FE 1: 88
22. *Rumex acetosella* — P; FE 1: 83

Chenopodiaceae

25. *Chenopodium murale* — A; FE 1: 94 27. *Salicornia fruticosa* — P; FAe: 123
26. *C. album* — A; FE 1: 94 28. *Salsola kali* — A; FE 1: 105

Amaranthaceae

29. *Amaranthus deflexus* — P; FE 30. *A. lividus* — A; FE 1: 110
1: 110

Portulacaceae

31. *Portulaca oleracea* — A; FE 1: 114

Caryophyllaceae

32. *Minuartia mediterranea* — A; 40. *Agrostemma githago* — A; FE
FE 1: 127 1: 157
33. *M. hamata* — A; FE 1: 127 41. *Silene italica* — P; FE 1: 163
34. *Stellaria media* — A; FE 1: 135 42. *S. vulgaris* subsp. *vulgaris* — P;
35. *Cerastium illyricum* — A; FE FE 1: 169
1: 142 43. *S. muscipula* — A; FE 1: 176
36. *Paronychia kapela* subsp. *chionea* 44. *S. behen* — A; FE 1: 177
— P; FE 1: 150 45. *S. gallica* — A; FE I: 179
37. *P. capitata* — P; FE 1: 150 46. *S. colorata* — A; FE 1: 180
38. *Herniaria hirsuta* — A; FE 47. *Petrorhagia armerioides* — P;
1: 152 FE 1: 187
39. *Spergularia rubra* — A; FE 48. *Dianthus diffusus* — P; FE
1: 155 1: 201

Ranunculaceae

49. *Nigella damascena* — A; FE 52. *Clematis cirrhosa* — P; FE
1: 210 1: 221
50. *Delphinium peregrinum* — A; 53. *Ranunculus muricatus* — A;
FE 1: 215 FE 1: 230
51. *Anemone coronaria* — P; FE 54. *R. millefoliatus* — P; FE 1: 231
1: 219 55. *R. sprunneranus* — P; FE 1: 231

Lauraceae

56. *Laurus nobilis* — P; FE 1: 246

Papaveraceae

57. *Papaver rhoeas* — A; FE 1: 248 60. *F. petteri* subsp. *thuretii* —
58. *Glaucium flavum* — B; FE 1: 250 A; FE 1: 257
59. *Fumaria macrocarpa* — A; FE 61. *F. officinalis* — A; FE 1: 258
1: 256

Capparidaceae

62. *Capparis spinosa* — P; FE 1: 259

Cruciferae

63. *Sisymbrium irio* — A; FE 1: 264 76. *Capsella bursa pastoris* — A;
 64. *Erysimum pusillum* subsp. *par-* FE 1: 316
nassii — P; FE 1: 272 77. *Aethionema saxatile* — A; FE
 65. *E. graecum* — B; FE 1: 272 1: 322
 66. *Malcolmia africana* — A; FE 78. *Biscutella didyma* — A; FE
 1: 277 1: 329
 67. *M. maritima* — A; FE 1: 277 79. *Cardaria draba* — P; FE 1: 333
 68. *M. graeca* — A; FE 1: 278 80. *Brassica cretica* — P; FE 1: 337
 69. *Matthiola sinuata* — B; FE 81. *B. tournefortii* — A; FE 1: 338
 1: 280 82. *Sinapis arvensis* — A; FE 1: 339
 70. *Cardamine graeca* — A; FE 83. *Eruca vesicaria* subsp. *sativa* —
 1: 289 A; FE 1: 340
 71. *Alyssum saxatile* — P; FE 1: 299 84. *Hirschfeldia incana* — A; FE
 72. *A. montanum* — P; FE 1: 301 1: 342
 73. *Clypeola jonthlaspi* — A; FE 85. *Cakile maritima* — A; FE 1: 343
 1: 307 86. *Raphanus raphanistrum* — A;
 74. *C. eriocarpa* — A; FE I: 307 FE 1: 346
 75. *Neslia paniculata* — A; FE 1: 315

Resedaceae

87. *Reseda alba* — A; FE 1: 347 88. *R. lutea* — A; FE 1: 348

Crassulaceae

89. *Umbilicus parviflorus* — P; FE 91. *S. tenuifolium* — P; FE 1: 359
 1: 351 92. *S. hispanicum* — A; FE 1: 363
 90. *Sedum sediforme* — P; FE 1: 358

Saxifragaceae

93. *Saxifraga hederacea* — A; FE 94. *S. tridactylites* — A; FE 1: 370
 1: 370

Platanaceae

95. *Platanus orientalis* — P; FE 1: 384

Rosaceae

96. *Rubus canescens* — P; FE 99. *Sarcopoterium spinosum* — P;
 2: 17 FE 2: 34
 97. *Rosa sempervirens* — P; FE 100. *Aphanes arvensis* — A; FE
 2: 27 2: 64
 98. *Sanguisorba minor* — P; FE 101. *Pyrus amygdaliformis* — P;
 2: 34 FE 2: 66
 102. *Prunus prostrata* — P; FE 2: 79

Leguminosae

103. *Cercis siliquastrum* — P; FE 2: 83
 104. *Ceratonia siliqua* — P; FE 2: 83
 105. *Parkinsonia aculeata* — P.
 106. *Anagyris foetida* — P; FE 2: 85
 107. *Calicotome villosa* — P; FE 2: 86
 108. *Genista acanthoclada* — P; FE 2: 100
 109. *Spartium junceum* — P; FE 2: 101
 110. *Lupinus angustifolius* — A; FE 2: 105
 111. *Robinia pseudacacia* — P; FE 2: 106
 112. *Colutea arborescens* — P; FE 2: 107
 113. *Astragalus sinaicus* — A; FE 2: 113
 114. *A. hellenicus* — P; FE 2: 117
 115. *A. graecus* — P; FE 2: 117
 116. *A. spruneri* — P; FE 2: 122
 117. *Psoralea bituminosa* — P; FE 2: 127
 118. *Vicia villosa* subsp. *microphylla* — A; FE 2: 132
 119. *V. tenuissima* — A; FE 2: 133
 120. *V. peregrina* — A; FE 2: 135
 121. *Lathyrus saxatilis* — A; FE 2: 141
 122. *L. sphaericus* — A; FE 2: 141
 123. *L. cicera* — A; FE 2: 142
 124. *L. aphaca* — A; FE 2: 143
 125. *Ononis ornithopodioides* — A; FE 2: 145
 126. *O. reclinata* — A; FE 2: 145
 127. *O. pubescens* — A; FE 2: 145
 128. *Melilotus italicica* — A; FE 2: 149
 129. *M. neapolitana* — A; FE 2: 149
 130. *Trigonella sprunerana* — A; FE 2: 151
 131. *T. coerulescens* — A; FE 2: 152
 132. *T. gladiata* — A; FE 2: 152
 133. *Medicago orbicularis* — A; FE 2: 155
 134. *M. marina* — P; FE 2: 156
 135. *M. rigidula* — A; FE 2: 156
 136. *M. littoralis* — A; FE 2: 156
 137. *M. arabica* — A; FE 2: 156
 138. *M. minima* — A; FE 2: 157
 139. *Trifolium tomentosum* — A; FE 2: 165
 140. *T. arvense* — A; FE 2: 167
 141. *T. stellatum* — A; FE 2: 168
 142. *T. angustifolium* — A; FE 2: 170
 143. *Dorycnium pentaphyllum* subsp. *herbaceum* — P; FE 2: 173
 144. *Lotus peregrinus* — P; FE 2: 176
 145. *Anthyllis hermanniae* — P; FE 2: 178
 146. *Coronilla emerus* subsp. *emerosides* — P; FE 2: 183
 147. *Hippocrepis unisiliquosa* — A; FE 2: 185
 148. *Onobrychis ebenoides* — P; FE 2: 188
 149. *O. caput-galli* — A; FE 2: 191

Oxalidaceae

150. *Oxalis corniculata* — P; FE 2: 192
 151. *O. pes-caprae* — P; FE 2: 193

Geraniaceae

152. *Geranium rotundifolium* — A; FE 2: 198
 153. *G. molle* — A; FE 2: 198
 154. *G. lucidum* — A; FE 2: 198

155. *Erodium laciniatum* — A; FE 2: 200
157. *E. gruinum* — A; FE 2: 201
156. *E. malacoides* — A; FE 2: 200
158. *E. cicutarium* — A; FE 2: 202

Zygophyllaceae

159. *Tribulus terrestris* — A; FE 2: 205

Linaceae

160. *Linum bienne* — B; FE 2: 209 161. *L. trigynum* — A; FE 2: 210

Euphorbiaceae

162. *Chrozophora tinctoria* — A; FE 2: 211
165. *E. helioscopia* — A; FE 2: 221
163. *Mercurialis annua* — A; FE 2: 212
166. *E. myrsinifolius* — P; FE 2: 221
167. *E. exigua* — A; FE 2: 222
168. *E. peplus* — A; FE 2: 222
164. *Euphorbia acanthothamnos* —
P; FE 2: 220

Rutaceae

169. *Ruta graveolens* — P; FE 2: 227

Simaroubaceae

170. *Ailanthus altissima* — P; FE 2: 231

Polygalaceae

171. *Polygala venulosa* — P; FE 2: 233

Anacardiaceae

172. *Pistacia terebinthus* — P; FE 2: 237
173. *P. lentiscus* — P; FE 2: 237
174. *Schinus molle* — P; FE 2: 237

Rhamnaceae

175. *Rhamnus lycioides* subsp. *graecus* — P; FE 2: 244

Malvaceae

176. *Malva cretica* — A; FE 2: 250 178. *Alcea rosea* — P; FE 2: 254
177. *M. sylvestris* — B; FE 2: 250

Thymelaeaceae

179. *Thymelaea hirsuta* — P; FE 2: 259 180. *T. tartonraira* — P; FE 2: 259

2: 259

Guttiferae

181. *Hypericum empetrifolium* — P; 183. *H. triquetrifolium* — P; FE
FE 2: 264 2: 269
182. *H. perfoliatum* — P; FE 2: 266

Cistaceae

184. *Cistus incanus* subsp. *creticus* — P; FE 2: 283
185. *C. parviflorus* — P; FE 2: 283
186. *C. monspeliensis* — P; FE 2: 283
187. *C. salvifolius* — P; FE 2: 284
188. *Tuberaria guttata* — P; FE 2: 286
189. *Helianthemum lavandulifolium* — P; FE 2: 287
190. *H. hymettium* — P; FE 2: 290
191. *Fumana arabica* — P; FE 2: 291
192. *F. thymifolia* — P; FE 2: 292

Tamaricaceae

193. *Tamarix gallica* — P; FE 2: 293

Cucurbitaceae

194. *Ecballium elaterium* — A; FE 2: 297
195. *Bryonia cretica* subsp. *dioica* — B; FE 2: 297

Cactaceae

196. *Opuntia ficus-indica* — P; FE 2: 300

Lythraceae

197. *Lythrum hyssopifolia* — A; FE 2: 301

Myrtaceae

198. *Eucalyptus robustus* — P; FE 2: 304
199. *E. globulus* — P; FE 2: 305

Punicaceae

200. *Punica granatum* — P; FE 2: 305

Araliaceae

201. *Hedera helix* — P; FE 2: 314

Umbelliferae

202. *Eryngium maritimum* — P; FE 2: 322
203. *E. campestre* — P; FE 2: 323
204. *Lagoetia cuminoides* — A; FE 2: 324
205. *Scandix pecten-veneris* — A; FE 2: 327
206. *Smyrnium olusatrum* — B; FE 2: 328
207. *S. perfoliatum* — B; FE 2: 328
208. *Foeniculum vulgare* — P; FE 2: 341
209. *Bupleurum fruticosum* — P; FE 2: 350
210. *Ammi majus* — A; FE 2: 353
211. *Ferulago nodosa* — P; FE 2: 359
212. *Tordylium apulum* — A; FE 2: 367
213. *Thapsia garganica* — P; FE 2: 370
214. *Daucus guttatus* — A; FE 2: 373
215. *D. carota* subsp. *maximus* — A; FE 2: 374

Ericaceae

216. *Erica arborea* — P; FE 3: 7 218. *Arbutus unedo* — P; FE 3: 11
 217. *E. manipuliflora* — P; FE 3: 7 219. *A. andrachne* — P; FE 3: 11

Primulaceae

Plumbaginaceae

Oleaceae

225. *Olea europaea* — P; FE 3: 55 226. *Phillyrea latifolia* — P; FE
 225a. *O. europaea* var *sylvestris* — 3: 55
 P; FE 3: 55

Apocynaceae

227. *Nerium oleander* — P; FE 3 : 68

Rubiaceae

228. *Asperula arvensis* — A; FE 4: 13
 229. *Galium incurvum* — P; FE 4: 26

230. *G. aparine* — A; FE 4: 35
 231. *G. murale* — A; FE 4: 36

Convolvulaceae

232. *Convolvulus boissieri* — P; FE 3: 80
 233. *C. dorycnium* — P; FE 3: 80
 234. *C. arvensis* — P; FE 3: 81
 235. *C. althaeoides* subsp. *althaeoides* — P; FE 3: 82
 235a. *C. althaeoides* subsp. *tenuissimus* — P; FE 3: 82

Boraginaceae

236. *Heliotropium europaeum* — A; FE 3: 85
 237. *Buglossoides arvensis* — A; FE 3: 37
 238. *Onosma frutescens* — P; FE 3: 91
 239. *Cerinthe minor* — A; FE 3: 94
 240. *Alkanna tinctoria* — P; FE 3: 96
 241. *Echium italicum* — B; FE 3: 98
 242. *E. platagineum* — A; FE 3: 99
 243. *E. parviflorum* — A; FE 3: 100
 244. *E. arenarium* — B; FE 3: 100
 245. *Anchusa undulata* — B; FE 3: 107
 246. *A. variegata* — A; FE 3: 109
 247. *Cynoglossum columnae* — A; FE 3: 120
 248. *C. creticum* — B; FE 3: 120

Verbenaceae

249. *Vitex agnus-castus* — P; FE 250. *Verbena officinalis* — P; FE
3: 122 3: 123

Labiatae

251. *Ajuga iva* — P; FE 3: 129 264. *Satureja thymbra* — P; FE
252. *Teucrium divaricatum* — P; 3: 164
FE 3: 132 265. *Calamintha nepeta* — P; FE
253. *T. polium* — P; FE 3: 134 3: 166
254. *Prasium major* — P; FE 3: 137 266. *Micromeria nervosa* — P; FE
255. *Marrubium vulgare* — P; FE 3: 169
3: 138 267. *M. juliana* — P; FE 3: 169
256. *Phlomis fruticosa* — P; FE 3: 145 268. *Origanum heracleoticum* — P;
FE 3: 171
257. *Lamium garganicum* — P; FE 3: 147 269. *Thymus capitatus* — P; FE
3: 174
258. *L. amplexicaule* — A; FE 3: 148 270. *Mentha pulegium* — P; FE
3: 184
259. *Ballota acetabulosa* — P; FE 3: 150 271. *Rosmarinus officinalis* — P;
FE 3: 187
260. *Stachys cretica* — P; FE 3: 153 272. *Salvia triloba* — P; FE 3: 189
261. *S. spruneri* — P; FE 3: 155 273. *S. pomifera* — P; FE 3: 190
262. *Glechoma hederacea* — P; FE 3: 161 274. *S. verbenaca* — P; FE 3: 192
275. *S. viridis* — P; FE 3: 192
263. *Melissa officinalis* — P; FE
3: 162

Solanaceae

276. *Hyoscyamus niger* — A; FE 278. *Datura stramonium* — A; FE
3: 195 3: 200
277. *Solanum nigrum* — A; FE 279. *Nicotiana glauca* — P; FE
3: 197 3: 201

Scrophulariaceae

280. *Verbascum undulatum* — B; 284. *Misopates orontium* — A; FE
FE 3: 213 3: 224
281. *V. sinuatum* — B; FE 3: 213 285. *Veronica glauca* — A; FE 2: 248
282. *Scrophularia peregrina* — A; 286. *V. arvensis* — A; FE 3: 249
FE 3: 218 287. *V. cymbalaria* — A; FE 3: 250
283. *S. heterophylla* — P; FE 3: 219 288. *Bellardia trixago* — A; FE
3: 269

Globulariaceae

289. *Globularia alypum* — P; FE 3: 283

Orobanchaceae

290. *Orobanche crenata* — A; FE 291. *O. pubescens* — P; FE 3: 290
3: 290

Plantaginaceae

292. *Plantago major* — P; FE 4: 39 296. *P. albicans* — P; FE 4: 43
293. *P. coronopus* — A; FE 4: 40 297. *P. bellardii* — A; FE 4: 43
294. *P. lanceolata* — P; FE 4: 42 298. *P. arenaria* — A; FE 4: 43
295. *P. lagopus* — P; FE 4: 43

Caprifoliaceae

299. *Lonicera implexa* — P; FE 4: 47 300. *L. etrusca* — P; FE 4: 47

Valerianaceae

301. *Centranthus ruber* — P; FE 302. *C. calcitrapae* — A; FE 4: 56
4: 55

Dipsacaceae

303. *Knautia integrifolia* — A; FE 304. *Scabiosa hymettia* — P; FE
4: 67 4: 69

Campanulaceae

305. *Campanula celsii* — B; FE 4: 81 307. *C. drabifolia* — A; FE 4: 88
306. *C. rupestris* — B; FE 4: 82

Compositae

308. *Bellis annua* — A; FE 4: 111 319. *A. chia* — A; FE 4: 155
309. *Conyza canadensis* — A; FE 320. *Chamomilla recutita* — A; FE
4: 120 4: 167
310. *Helichrysum stoechas* subsp. 321. *Chrysanthemum segetum* — A;
barrelieri — P; FE 4: 129 FE 4: 168
311. *Phagnalon graecum* — P; FE 322. *C. coronarium* — A; FE 4: 169
4: 133 323. *Senecio vulgaris* — A; FE 4: 204
312. *Dittrichia viscosa* — P; FE 324. *Calendula arvensis* — A; FE
4: 137 4: 207
313. *D. graveolens* — A; FE 4: 137 325. *Carlina corymbosa* subsp. *graeca*
314. *Pulicaria odora* — P; FE 4: 137 — B; FE 4: 209
315. *Pallenis spinosa* — A; FE 326. *Atractylis cancellata* — A; FE
4: 139 4: 211
316. *Xanthium strumarium* — A; 327. *Echinops sphaerocephalus*
FE 4: 143 subsp. *albidus* — P; FE 4: 214
317. *X. spinosum* — A; FE 4: 143 328. *E. microcephalus* — P; FE
318. *Anthemis tomentosa* — A; FE 4: 214
4: 154 329. *Carduus pycnocephalus* — A;
FE 4: 231

330. *Cirsium creticum* — P; FE 4: 242
331. *Picnomon acarna* — A; FE 4: 242
332. *Notobasis syriaca* — A; FE 4: 242
333. *Ptilostemon chamaepeuce* — P; FE 4: 243
334. *Onopordon illyricum* — B; FE 4: 247
335. *Silybum marianum* — A; FE 4: 249
336. *Centaurea raphanina* subsp. *mixta* — P; FE 4: 269
337. *C. attica* — P; FE 4: 275
338. *C. spinosa* subsp. *tomentosa* — P; FE 4: 282
339. *C. solstitialis* — B; FE 4: 284
340. *Crupina crupinastrum* — A; FE 4: 301
341. *Carthamus lanatus* — A; FE 4: 303
342. *Scolymus hispanicus* — B; FE 4: 304
343. *Cichorium intybus* — P; FE 4: 304
344. *Rhagadiolus stellatus* — A; FE 4: 308
345. *Picris pauciflora* — A; FE 4: 317
346. *Scorzoneroides crocifolia* — P; FE 4: 320
347. *S. lanata* — P; FE 4: 322
348. *Tragopogon porrifolius* — B; FE 4: 323
349. *Reichardia picroides* — P; FE 4: 325
350. *Sonchus oleraceus* — A; FE 4: 327
351. *Steptorhamphus tuberosus* — P; FE 4: 328
352. *Lactuca serriola* — A; FE 4: 330
353. *Taraxacum megalorhizon* — P; FE 4: 334
354. *T. officinale* — P; FE 4: 342
355. *Chondrilla ramosissima* — B; FE 4: 343
356. *Crepis foetida* — A; FE 4: 354
357. *C. micrantha* — A; FE 4: 355

b. *Monocotyledones*

Liliaceae

358. *Merendera attica* — P; GF A: 79
359. *Colchicum pusillum* — P; FAe: 709
360. *C. latifolium* — P; FAe: 709
361. *Asphodelus microcarpus* — P; FAe: 710
362. *Gagea peduncularis* — P; FAe: 712
363. *G. arvensis* — P; FAe: 712
364. *Allium ampeloprasum* — P; FAe: 715
365. *A. neapolitanum* — P; FAe: 718
366. *A. roseum* — P; FAe: 718
367. *A. stamineum* — P; FAe: 718
368. *Fritillaria graeca* — P; FAe: 721
369. *Lloydia graeca* — P; FAe: 722
370. *Drimia maritima* — P; STEARN 1978
371. *Scilla autumnalis* — P; FAe: 724
372. *Ornithogalum atticum* — P; FAe: 725
373. *O. tenuifolium* — P; FAe: 725
374. *O. umbellatum* — P; FAe: 725
375. *O. nanum* — P; FAe: 726
376. *Bellevalia hyacinthoides* — P; PERSSON & WENDELBO 1979
377. *Leopoldia comosa* — P; BENTZER 1973: 75
378. *Muscari commutatum* — P; FAe: 731

379. *Asparagus acutifolius* — P; FAe: 732
 380. *A. aphyllus* — P; F733 Ae: 381. *Ruscus aculeatus* — P; FAe:
 733
 380. *A. aphyllus* — P; F733 Ae: 382. *Smilax aspera* — P; FAe: 734
Amaryllidaceae
383. *Stenbergia sicula* — P; FAe: 735
Agavaceae
384. *Agave americana* — P; FAe: 736
Iridaceae
385. *Crocus laevigatus* — P; FAe: 738
 386. *C. sieberi* — P; FAe: 738
 387. *C. cartwrightianus* — P; FAe: 738
 388. *C. cancellatus* — P; FAe: 738
Juncaceae
392. *Juncus compressus* — P; FAe: 745
 393. *J. maritimus* — P; FAe: 746
 394. *Luzula nodulosa* — P; FAe: 747
Cyperaceae
395. *Carex distachya* — P; FAe: 754
 396. *C. hispida* — P; FAe: 755
Gramineae
397. *Bromus madritensis* — A; FAe: 759
 398. *B. rubens* — A; FAe: 759
 399. *B. sterilis* — A; FAe: 760
 400. *B. intermedius* — A; FAe: 761
 401. *Agropyron repens* — P; FAe: 764
 402. *Aegilops ovata* — A; FAe: 766
 403. *Ae. triaristata* — A; FAe: 766
 404. *Hordeum crinitum* — A; FAe: 768
 405. *H. murinum* — A; FAe: 769
 406. *Arundo donax* — P; FAe: 771
 407. *Briza maxima* — A; FAe: 774
 408. *B. spicata* — A; FAe: 775
 409. *Dactylis glomerata* — P; FAe: 775
 410. *Cynosurus echinatus* — P; FAe: 777
 411. *Poa annua* — A; FAe: 778
 412. *P. bulbosa* — P; FAe: 779
 413. *Scleropoa rigida* — A; FAe: 786
 414. *Lolium temulentum* — A; FAe: 787
 415. *L. perenne* — P; FAe: 788
 416. *Koeleria phleoides* — A; FAe: 790
 417. *Avena sterilis* — A; FAe: 792
 418. *A. barbata* — A; FAe: 792
 419. *Aira capillaris* — A; FAe: 794
 420. *Lagurus ovatus* — A; FAe: 797
 421. *Stipa bromoides* — P; FAe: 801
 422. *S. lagascae* — P; FAe: 802
 423. *S. tortilis* — A; FAe: 802
 424. *Oryzopsis miliacea* — P; FAe: 802
 425. *O. coerulescens* — P; FAe: 803
 426. *Digitaria sanguinalis* — A; FAe: 806
 427. *Cynodon dactylon* — P; FAe: 806
 428. *Panicum crus-galli* — A; FAe: 806
 429. *Setaria viridis* — A; FAe: 807

Orchidaceae

- | | |
|--|---|
| 430. <i>Ophrys apifera</i> — P; FAe: 811 | 435. <i>S. lingua</i> — P; FAe: 824 |
| 431. <i>O. fusca</i> — P; FAe: 815 | 436. <i>Anacamptis pyramidalis</i> — P; |
| 432. <i>O. heldreichii</i> — P; FAe: 817 | FAe: 828 |
| 433. <i>O. lutea</i> — P; FAe: 817 | 437. <i>Neotinea intacta</i> — P; FAe: 830 |
| 434. <i>Serapias laxiflora</i> — P; FAe: | 438. <i>Orchis laxiflora</i> — P; GF A: 230 |
| 822 | 439. <i>O. quadripunctata</i> — P; FAe: |
| | 837 |

Araceae

- | | |
|---|--|
| 440. <i>Arum italicum</i> — P; FAe: 844 | 442. <i>Biarum spruneri</i> — P; GF A: |
| 441. <i>Dracunculus vulgaris</i> — P; | 242 |
| FAe: 845 | 443. <i>Arisarum vulgare</i> — P; FAe: |
| | 845 |

References

- ARRIGONI E. 1971. Elements for the representation of the Attica landscape at the classic era. — Athens. (in Greek).
- BENTZER B. 1973. Taxonomy, variation and evolution in representatives of *Leopoldia* PARL. (*Liliaceae*) in the Southern and Central Aegean. — Bot. Notiser 126 (1): 69—132.
- DIAPOLIS Ch. 1939—1949. Greek Flora. I—III. — Athens. (in Greek).
— 1958. From the flora of Parnis. “To Vouno” 1958 (Nov.—Dec.): 163—188. — Athens. (in Greek).
— 1968. Report about the Botanical Garden “Julia and Alex Diomidous”. — Athens. (Unpublished). (in Greek).
- LEPSIUS R. 1893. Geologie von Attica. — Berlin.
- PERSSON K. & WENDELBO P. 1979. *Bellevalia hyacinthoides*, a new name for *Strangweja spicata* (*Liliaceae*). — Bot. Notiser 132 (1): 65—70.
- RECHINGER K. H. fil. 1943. Flora Aegaea. — Denkschr. Akad. Wiss. Wien, math.-naturw. Kl. 105 (1): 1—924.
- STEARNS W. T. 1978. Mediterranean and Indian species of *Drimia* (*Liliaceae*): a nomenclatural survey with special reference to the medicinal squill, *D. maritima* (syn. *Urginea maritima*). — Ann. Mus. Goulandris 4: 199—210.
- TATARIS A. 1967. Remarks on the structure of the Skaramanga-Egaleo Mountain—Piraeus—Athens. (Attica area). — Bull. hellen. geolog. Comp. 1: 52—67. Athens. (in Greek).
- TUTIN T. & al. (Eds.). 1964—1976. Flora Europaea. 1—4. — Cambridge University Press.
- ZERLENTIS K. 1959. The plants of Lyceabettus. — Athens. (in Greek).
— 1965. Contribution to the flora of Mount Hymettus. — Athens. (in Greek).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Phyton, Annales Rei Botanicae, Horn](#)

Jahr/Year: 1980

Band/Volume: [20_3_4](#)

Autor(en)/Author(s): Sarlis George P.

Artikel/Article: [The Flora of Egalo \(Attica, Greece\). 261-278](#)